

CASE REPORT

PEER REVIEWED | OPEN ACCESS

Toxic megacolon and perforation of the right colon due to sigmoid stenosis associated to chronic diverticulitis

Maria Tudela Lerma, Ana Moreno Hidalgo, Benjamin Diaz Zorita

ABSTRACT

Introduction: Diverticulosis is a common benign disease in the population over the age of 60. It ranges from asymptomatic to complicate with bleeding or perforation. "Chronic diverticulitis" is defined by its chronic clinical course and luminal obstructive change (stenosis), which may rarely lead in large bowel obstruction. Colonic dilation can cause toxic megacolon and perforation. **Case Report:** This is a rare case of toxic megacolon accompanied by perforation of the right colon due to chronic dilatation caused by stenosis of the sigmoid colon as a complication of diverticulitis. The patient consulted in emergency for abdominal pain and developed severe septic shock. The computed tomography showed dilatation of the colon with perforation and large retroperitoneal abscess. An emergency total colectomy was performed. **Conclusion:** To the best of our knowledge there are very few reports in the literature referring to toxic megacolon and perforation of the right colon due to stenosis of the sigmoid colon as a result of chronic diverticulitis.

Keywords: Colonic dilatation, Colon perforation, Colon stenosis, Diverticulitis complication, Toxic megacolon

How to cite this article

Lerma MT, Hidalgo AM, Zorita BD. Toxic megacolon and perforation of the right colon due to sigmoid stenosis associated to chronic diverticulitis. J Case Rep Images Surg 2018;4:100056Z12ML2018.

Article ID: 100056Z12ML2018

doi: 10.5348/100056Z12ML2018CR

INTRODUCTION

Diverticulosis is a common benign disease in the population over the age of 60. It affects the sigmoid and descending colon in more than 90% of patients [1, 2].

The spectrum of diverticular disease is wide, covering different clinical scenarios; it ranges from asymptomatic to different complications [3].

Recently a new concept of diverticulitis was proposed: "chronic diverticulitis" [4, 5] it is defined by its chronic clinical course and luminal obstructive change (stenosis). This disease is a different pathologic entity considered by the common development of chronic obstructive symptoms. The chronic inflammation and recurrent diverticulitis in the left-sided colon can cause fibrosis subsequent stenosis and finally obstruction of the colon. The symptoms include vomiting, constipation and abdominal distension. Compared with acute diverticulitis it is characterized by a lack of abdominal pain, tenderness and fever [6].

Acute diverticulitis is the second cause of large bowel obstruction due to a spasm, anearby abscess or a fibrotic scar; however, chronic diverticulitis (stenosis) is a rare

Maria Tudela Lerma¹, Ana Moreno Hidalgo², Benjamin Diaz Zorita³

Affiliations: ¹PhD, General Surgeon, Esophagogastric Department Gregorio Marañon University General Hospital, Madrid, Spain; ²PhD, General Surgeon, Gregorio Marañon University General Hospital, Madrid, Spain; ³PhD, General Surgeon, Hepatobiliary Department, Gregorio Marañon University General Hospital, Madrid, Spain.

Corresponding Author: Maria Tudela Lerma, Duque de Ses- to 17, 5° C, 28009, Madrid, Spain; Email: matuler@hotmail.com

Received: 16 August 2018

Accepted: 13 September 2018

Published: 11 October 2018

cause of obstruction. It is supposed to be the cause of only 10% of large bowel obstructions [7]. Nevertheless if the stenosis is severe, a progressive dilation of the proximal colon (cecum >9 cm or transverse colon >6 cm) may lead to toxic megacolon followed by colonic ischemia-necrosis and perforation [8].

This report present and discusses a case of sigmoid stenosis due to diverticulitis and chronic dilation of the colon, which evolved in progressive proximal colon dilatation complicated by toxic megacolon and perforation of the proximal colon.

CASE REPORT

A 81-year-old women with medical history of two episodes of acute diverticulitis with pararectal abscess managed conservatively with antibiotic therapy and percutaneous drainage. Along the last year the patient presented chronic symptoms vomiting, constipation and abdominal distension. A colonoscopy revealed a partial inflammatory sigma stenosis; malignancy was discarded with two biopsies.

The patient consulted in emergency department with severe diffuse abdominal pain, which had started 24 hours earlier. She besides reported fever and vomiting.

At physical exploration she was with poor general condition, hypotensive 95 /64 mmHg, tachycardic with 115 beats/min and the temperature was 38°C. Abdominal examination revealed a distended, tympanic, diffusely painful abdomen with tenderness and abdominal guarding.

The total leukocyte count at admission was 4.100/mm³ with 89% neutrophils, increased acute phase reactants, impaired renal function with creatinine 1.36 mg/dL and high blood lactic acid. A CT- scan showed located complete stenosis of the sigmoid with dilation of the proximal colon. Discarded pneumomediastinum, pneumoperitoneum, retroperitoneum and a large collection in right hemiabdomen Figure 1(A–C). The above clinical and laboratory findings were compatible with toxic megacolon.

The patient was taken to the operating room where laparotomy revealed fecaloid peritonitis of the four quadrants, due to 5 cm perforation of the right colon (Figure 2) with dilatation of the colon due to the stenosis of the sigma. A total colectomy with terminal ileostomy was performed. Biopsy revealed colonic ischemia and diverticulosis of the colon as well as deformity of the sigma wall by fibrosis and severe luminal narrowing of 7 cm Figure 3(A–C).

DISCUSSION

Diverticulosis is a common anatomical condition, which appears to be age dependent. Recent studies have shown that the presence of diverticula represents the




Figure 1(A–C): Abdominal CT showed: (A, B) Pneumoperitoneum, the orifice of the right colon and the large abscess of 14x10x18 cm. Stenotic lesion of the sigmoid colon (C).




Figure 2: Operating findings: fecaloid peritonitis due to right colon perforation.

most common non-neoplastic finding during screening colonoscopy [9]. Among patients with diverticulosis, 15–25% are expected develop diverticulitis in their lifetime [10], the incidence of acute diverticulitis increases with age, especially over the age of 50 [11, 12], although a recent study suggests that this proportion may be much lower [13].

It may be difficult to distinguish “chronic diverticulitis” from other diseases, which are complicated by intestinal stenosis including cancer, ulcerative colitis, Crohn’s disease, and ischemic colitis. The diagnostic can be




Figure 3(A–C): Resected specimen. Total colectomy (A). Right colon perforation, length 5 cm (B). Stenosis of the sigmoid colon, length 7 cm (C).

settled with the clinical characteristics (symptoms of ileus and past history with diverticulitis) and the radiographic findings. For imaging, it is useful a barium enema, computerized tomography, and endoscopy [5].

The differential diagnosis of sigmoid stenosis secondary to diverticulitis from cancer could be difficult and is a challenge for surgeons. CT colonography is an excellent technique to distinguish between both etiologies. It can evaluate the proximal colon; however, it is also contraindicated for evaluating acute abdominal conditions, like diverticulitis or the acute phase of inflammatory bowel disease, because of potential complications of air insufflation [14].

The sigmoid's stricture can cause progressive distension of the colon and increase the intraluminal pressure, which may provoke ischemia of all bowel wall layers and finally necrosis and perforation.

Toxic megacolon is a rare but severe and potentially fatal complication of colonic inflammation with high morbidity and mortality, and surgical intervention is necessary in up to 80% of cases. It is most commonly considered a complication of inflammatory bowel disease and less frequent of ischemic or infective colitis. Chronic sigma stenosis due to diverticulitis is an exceptional cause of megacolon.

The consistent feature of toxic megacolon is the radiographic evidence of total or segmental colonic

distension of >6 cm. In comparison to other aetiologies of colonic dilatation, it is defined by the additional presence of systemic toxicity and inflammatory or infectious aetiology of the underlying disease. The most commonly used clinical criteria for the diagnosis of toxic megacolon include three of the four following main criteria: Fever, tachycardia, leukocytosis, or anemia. In addition, one of the following criteria should also be met: Dehydration, altered level of consciousness, electrolyte imbalance, or hypotension [15]. The patient in the case was in septic shock.

Presently, the timing of surgery in toxic megacolon remains controversial. Avoiding the necessity of surgery is the goal for all medical treatments; however, delaying surgical therapy can increase the risk of complications leading to a poor prognosis [16].

On the other hand it is debated if the prophylactic surgery of the chronic diverticulitis is indicated to avoid these severe complications. Today, the number of episodes alone is no longer regarded as a conclusive indication for surgery. It depends on the individual case and takes risk factors, complications, age, severity of episode, as well as the patient's personal circumstances and comorbidities. Chronic recurrent, uncomplicated diverticulitis should only be operated on after a careful risk-benefit analysis during an inflammation-free interval [17].

CONCLUSION

Recurrent episodes of diverticulitis, which may be subclinical, can initiate progressive fibrosis and stricturing of the colonic wall in the absence of ongoing inflammation. Therefore, it can cause large bowel obstruction and finally toxic megacolon. The policy of prophylactic surgery, the timing and the appropriateness of treatment of sigmoid diverticulitis remain a topic of controversy.

REFERENCES

1. Stollman NH, Raskin JB. Diverticular disease of the colon. *J Clin Gastroenterol* 1999 Oct;29(3):241–52.
2. Morris AM, Regenbogen SE, Hardiman KM, Hendren S. Sigmoid diverticulitis: A systematic review. *JAMA* 2014 Jan 15;311(3):287–97.
3. Feuerstein JD, Falchuk KR. Diverticulosis and diverticulitis. *Mayo Clin Proc* 2016 Aug;91(8):1094–104.
4. Strate LL, Modi R, Cohen E, Spiegel BM. Diverticular disease as a chronic illness: Evolving epidemiologic and clinical insights. *Am J Gastroenterol* 2012 Oct;107(10):1486–93.
5. Sheiman L, Levine MS, Levin AA, et al. Chronic diverticulitis: Clinical, radiographic, and pathologic findings. *AJR Am J Roentgenol* 2008 Aug;191(2):522–8.
6. Miyamoto R, Koizumi M, Terashima T, et al. Chronic diverticulitis of the large intestine characterized by stenosis symptoms in Japan: Report of 6 cases. *Clin J Gastroenterol* 2012 Feb;5(1):47–52.

7. Buckley O, Geoghegan T, O'Riordain DS, Lyburn ID, Torreggiani WC. Computed tomography in the imaging of colonic diverticulitis. Clin Radiol 2004 Nov;59(11):977–83.
8. Antonopoulos P, Almyroudi M, Kolonia V, Kouris S, Troumpoukis N, Economou N. Toxic megacolon and acute ischemia of the colon due to sigmoid stenosis related to diverticulitis. Case Rep Gastroenterol 2013 Sep 11;7(3):409–13.
9. Scarpignato C, Barbara G, Lanis A, Strate LL. Management of colonic diverticular disease in the third millennium: Highlights from a symposium held during the United European Gastroenterology Week 2017. Therap Adv Gastroenterol 2018 May 20;11:1756284818771305.
10. Bevan R, Lee TJ, Nickerson C, et al. Non-neoplastic findings at colonoscopy after positive faecal occult blood testing: Data from the english bowel cancer screening programme. J Med Screen 2014 Jun;21(2):89–94.
11. Stollman N, Raskin JB. Diverticular disease of the colon. Lancet 2004 Feb 21;363(9409):631–9.
12. Munie ST, Nalamati SPM. Epidemiology and pathophysiology of diverticular disease. Clin Colon Rectal Surg 2018 Jul;31(4):209–13.
13. Shahedi K, Fuller G, Bolus R, et al. Long-term risk of acute diverticulitis among patients with incidental diverticulosis found during colonoscopy. Clin Gastroenterol Hepatol 2013 Dec;11(12):1609–13.
14. Ripollés T, Martínez-Pérez MJ, Gómez Valencia DP, Vizuite J, Martín G. Sigmoid stenosis caused by diverticulitis vs. carcinoma: Usefulness of sonographic features for their differentiation in the emergency setting. Abdom Imaging 2015 Oct;40(7):2219–31.
15. Ramanathan S, Ojili V, Vassa R, Nagar A. Large bowel obstruction in the emergency department: Imaging spectrum of common and uncommon causes. J Clin Imaging Sci 2017 Apr 5;7:15.
16. Carabotti M, Annibale B. Treatment of diverticular disease: An update on latest evidence and clinical implications. Drugs Context 2018 Mar 21;7:212526.
17. Jurowich CF, Germer CT. Elective surgery for sigmoid diverticulitis - indications, techniques, and results. Viszeralmedizin 2015 Apr;31(2):112–6.

Author Contributions

Maria Tudela Lerma – Substantial contributions to conception and design, Acquisition of data, Analysis and interpretation of data, Drafting the article, Revising it critically for important intellectual content, Final approval of the version to be published

Ana Moreno Hidalgo – Substantial contributions to conception and design, Acquisition of data, Analysis and interpretation of data, Drafting the article, Revising it critically for important intellectual content, Final approval of the version to be published

Benjamin Diaz Zorita – Substantial contributions to conception and design, Acquisition of data, Analysis and interpretation of data, Drafting the article, Revising it critically for important intellectual content, Final approval of the version to be published

Guarantor of Submission

The corresponding author is the guarantor of submission.

Source of Support

None.

Consent Statement

Written informed consent was obtained from the patient for publication of this case report.

Conflict of Interest

Authors declare no conflict of interest.

Data Availability

All relevant data are within the paper and its Supporting Information files.

Copyright

© 2018 Maria Tudela Lerma et al. This article is distributed under the terms of Creative Commons Attribution License which permits unrestricted use, distribution and reproduction in any medium provided the original author(s) and original publisher are properly credited. Please see the copyright policy on the journal website for more information.

Access full text article on
other devices



Access PDF of article on
other devices





INTERNATIONAL JOURNAL OF
CASE REPORTS AND IMAGES



VIDEO JOURNAL OF
CLINICAL RESEARCH



VIDEO JOURNAL OF
BIOMEDICAL SCIENCE




INTERNATIONAL JOURNAL OF
HEPATOBIILIARY AND
PANCREATIC DISEASES



INTERNATIONAL JOURNAL OF
BLOOD TRANSFUSION AND
IMMUNOHEMATOLOGY



EDORIUM JOURNAL OF
OPHTHALMOLOGY



EDORIUM JOURNAL OF
MEDICINE



EDORIUM JOURNAL OF
CARDIOTHORACIC AND
VASCULAR SURGERY



JOURNAL OF CASE REPORTS
AND IMAGES IN ORTHOPEDICS
AND RHEUMATOLOGY



EDORIUM JOURNAL OF
PSYCHOLOGY



EDORIUM JOURNAL OF
CELL BIOLOGY



JOURNAL OF CASE REPORTS AND IMAGES IN
DENTISTRY




EDORIUM JOURNAL OF
CANCER




EDORIUM JOURNAL OF
PSYCHIATRY



JOURNAL OF CASE REPORTS AND
IMAGES IN INFECTIOUS DISEASES



EDORIUM JOURNAL OF
ANATOMY AND EMBRYOLOGY



EDORIUM JOURNAL OF
SURGERY



JOURNAL OF CASE REPORTS
AND IMAGES IN PATHOLOGY



EDORIUM JOURNAL OF
ANESTHESIA